



<image><image><image><image><image><image><image><image><text><text><text>









9

Disclosures

• No financial relationships to disclose

Prevention of ASCVD in South Asians Impact of Diet Modification and Physical Activity in Primary Prevention

Geeta Sikand, MA, RDN, FAND, CDE, CLS, FNLA Associate Clinical Professor of Medicine (Cardiology) Director of Nutrition Univ of Calif Irvine Preventive Cardiology Program University of California, Irvine

10

Objectives

1. Enumerate the impact of food acculturation on the dietary patterns of South Asians (SA) and associated ASCVD risk factors.

2. Explain the impact of diet and lifestyle modification on ASCVD risk factors.

3. Describe SA diet and lifestyle goals and resources to help practitioners implement culturally tailored AHA/ACC recommended heart healthy dietary patterns.



WHO Facts: South Asians

- CHD strikes South Asians at an *earlier age* (almost 33% earlier) and with higher mortality rates than other demographics.
- 50% of all heart attacks in Indian men occur under 50 years of age and 25% of all heart attacks occur under 40 years of age.
- India accounts for approximately 60% of the world's heart disease burden, despite less than 20% of the world's population.
- India: World's capital for diabetes e.g. in Hyderabad, India, 20% of the entire adult population is diabetic.

WHO, 2011; Chiu M et al., Diabetes Care, 2011

14

ACC/AHA 2018 Cholesterol Management Guidelines: Risk-Enhancing Factors

- Family history of premature ASCVD
- Primary hypercholesterolemia
- Metabolic syndrome
- Chronic kidney disease
- Chronic inflammatory conditions
- Premature menopause
- High-risk race/ethnicities (e.g., South Asian ancestry)
- Lipid/biomarkers
- Elevated high-sensitivity C-reactive protein
- Elevated Lp(a)
 - Grundy et al. Circulation 2018


ASCVD Risk Factors in South Asians

Relatively lower BMI but more


 The lower BMI cut-point for over weight in South Asians was lowered from 24.9 to 23.0 due to early

central abdominal obesity.

15







High diabetes rate.

insulin resistance.

High risk of early heart disease.

Grundy Circulation 2018, Vogelman Circulation 2018, Jacobson J

MASALA and MESA on Pooled Risk Equation

 Discordance between 10-year cardiovascular risk estimates using the ACC/ AHA 2013 estimator and coronary artery calcium in SA individuals was noted when compared with 5 racial/ethnic groups when Comparing MASALA and MESA studies

Al Rifai, M., Atherosclerosis, https://doi.org/10.1016/j.atherosclerosis.2018.09.015

19

Impact of Acculturation in South Asians' dietary patterns and ASCVD Risk

- Increased intake of animal protein, fried snacks, sweets & high-fat dairy: Increased insulin resistance and reduced HDL-C.
- Adhere to a traditional plus western dietary pattern: Increased obesity and hypertension.
- Adhere to a western dietary pattern: Increased risk for MetS.

Eriksen et al. PLoS One. 2015

21



Vegetarian vs. Non-Vegetarians Dietary **Patterns** Vegetarians (Lacto) Non-vegetarians Obesity Dyslipidemia Dyslipidemia Overweight ASCVD Consume excessive Consume excessive animal products. calories, refined carbs, saturated fats from high fat dairy (ghee, butter, whole milk, cream) and coconut oil. Eriksen et al. PLoSOne. 2015

Impact of Acculturation on Incident CVD in South Asians

2- to 3-fold higher risk of incident CVD in non-adherers to heart healthy behaviors e.g.

- Fewer fruits and vegetables
- Sedentary lifestyle
- Smokers
- Excessive alcohol use

Parackal et al. Curr. Diabetes Rev. 2017

20

SA Dietary Patterns vary by Region and Religion

However, all consume high amounts of saturated fat & refined carbs:

- Ghee, butter, whole milk, cream
- Shortening, fried foods, coconut oil
- Potatoes, white rice, pizza
- Repeat use of cooking oil in deep fried savory snacks
- Sugar sweetened beverages

Parackal Curr. Diabetes Rev. 2017

The Mediators of Atherosclerosis in South Asians Living in America (MASALA) Study

 Kanaya et al. Acculturation & Subclinical Atherosclerosis among U.S. South Asians: Findings from the MASALA study. J Clin Exp Res Cardiol 2014.

MASALA Study Design Kanaya et al. 2014 MASAL MASALA **MESA** Ages 45-84 years Ages 40-84 years N = 900 N = 6,500 Only South 4 ethnic Asians groups Two sites (UCSF and NWU) 6 sites (Columbia, Hopkins, NWU, Minnesota, Pilot study (n=150; 2006-2007) UCLA, Wake Forest) Oct 2010-March 2013 Started in 2000: Exam 5, 2010-2012


26

25




Kanaya et al. Acculturation & Subclinical Atherosclerosis among U.S. South Asians: Findings from the MASALA study. J Clin Exp Res Cardiol 2014.

27










diabetes, stroke, MetS

31



33

Summary of Changes in SA Dietary Patterns upon Acculturation in the West 1. Decrease in beans, lentils, fruits & vegetables. 2. Increase in potatoes, dairy, oil, meat & fish. **3**. Increase in fast foods due to increased frequency of eating out. 4. Increase in rich traditional foods e.g. refined carbs, sweets and snacks prepared with ghee, coconut oil, butter and shortening. 5. Increase in western desserts and snacks.

32

Impact of Low Glycemic Load **Carbs: Vegetables and Fruits** "Global Burden of Disease Study 2010" showed low fruit and vegetable consumption was the top risk factor for causing greatest loss of health

Eriksen PLoS One. 2015



Lozano R et al. Lancet 2012

worldwide.

34

36



Fruits and Veggies are rich in **Flavanoids**

- Reduce oxidative stress
- Anti-inflammatory
- Prevent thrombus formation
- Improve Endothelium funx
- Improve lipids, BP & glycemia
- Sikand, Kris-Etherton et al. Current Cardiology Reports February 2015.







Low Glycemic Load Carbs Decrease • **V** CAD risk 20-30% inflammation ● High Fiber/ → markers promote satiety Less Insulin resistance - **↓** FBS & A1c Improved Lipid profiles —**↓** TG's, **↑** HDL Sikand, Kris-Etherton et al. Current Cardiology Reports 2015

Objective 2

Describe the impact of diet and lifestyle

modification on ASCVD risk factors in SA.


Grundy Circulation 2018, Vogelman Circulation 2018, Jacobson J Clin. Lipidol. 2015

 ...little population-based data in U.S. (some in UK and Australia), and no


longitudinal studies worldwide.

38

37



39



- 16 wks. nonrandomized trial (Chicago)
- Women (pre-T2D)
 Intervention: Healthy diet plus
- twice weekly exercise intervention Improved: BMI
- Kandula et al. *Transl J Am Coll* Sports Med. 2016

- 8 wks. RCT (SFO Bay area)
- (CURE-D study).
- Women with T2D
 Intervention: Twice weekly culturally
- relevant exercise (Bollywood dancing).
 Improved: BMI and A1c
- Natesan et al. *BMJ* Open Diabetes Res. Care. 2015

South Asian Heart Lifestyle Intervention (SAHELI) Study Chicago

- 6 mos. intervention with focus on a healthy diet in group sessions.
- Improved: weight status, A1c, physical activity and stress management skills.

Kandula et al. BMC Public Health 2015

40

Diet and Ex Intervention in SA with Pre-diabetes

- 3 mos. SFO Bay Area
- Overweight women with insulin resistance.
- Intervention: Low calorie, relatively low carb dietary pattern.
- Improved: BMI, insulin sensitivity & CVD risk factors.
- Backes AC et al. Asia Pac J Clin Res Nutr. 2008

43

6 mos. NY community based

- Adults with prediabetes.
- Intervention: Healthy diet, exercise & stress management.
- Improved: BMI, BP, BG, total-C, food habits and exercise.

Islam NS et al. Int J Environ Res Public Health. 2014

2019 AHA/ACC Prevention Guidelines for Adults

Eat a heart-healthy **dietary pattern**

- Emphasize plant-based foods, lean protein and fish.
- Limit foods high in saturated fats and dietary cholesterol.
- Minimize trans fat, sodium (salt), processed meats, refined carbs and sweetened beverages.

Be **physically active** most days of the week, 150 min/wk moderate-intensity e.g. brisk walking or 75 min/wk for high intensity e.g. jogging.

Arnott et al. Circulation 2019

45

Culturally Relevant Resources from Academy of Nutrition and Dietetics

Indians in Nutrition and Dietetics Member Interest Group of the Academy of Nutrition and Dietetics

- "Ready, Set, Start Counting" A carbohydrate counting tool for managing your diabetes for Asian Indians.
- Indian diet-healthy plate.
- Find a RD by zip code <u>www.eatright.org</u>



Objective 3

 List diet and lifestyle goals and practical resources to help practitioners implement culturally relevant AHA/ACC recommended heart healthy dietary patterns in SA patients.

44


US and Indian ASCVD Prevention Guidelines Recommend

Same guiding principles for a heart healthy diet apply to SA.

Implementation will differ based on cultural/ethnic and personal food preferences.

- DASH (Dietary Approaches to Stop Hypertension) dietary pattern
- Healthy Mediterranean-style dietary pattern (Healthy US dietary pattern)
- Healthy Vegetarian/Vegan dietary pattern.

Jacobson 2015, Vanhorn 2016, Iyengar 2016, Grundy 2018, Arnett 2019



Recommended Dietary Patterns to Achieve Adherence to the AHA/ACC Guidelines Van Horn et al. AHA Scientific Statement *Circulation 2016*

- Evidence-based dietary recommendations to facilitate adherence to the recent AHA/ACC Prevention Guidelines, 2015–2020 Dietary Guidelines for Americans to achieve AHA's 2020 Strategic Impact Goals.
- Provide guidance for achieving adherence to a heart-healthy dietary pattern and accommodate personal, cultural and ethnic food choices.

2016 AHA Scientific Statement Recommendations for Populations Worldwide

- Populations worldwide should follow the same guiding principles for a heart healthy dietary pattern to achieve the AHA 2020 goals.
- Implementation should be based on personal, socio-economic, cultural, ethnic and regional food preferences.
- Enhance acculturation by substituting heart-healthy ingredients in recipes for traditional foods.

Vanhorn et al. AHA Scientific Statement. Circulation 2016

49

Nutrition goals for optimizing lipids and reducing ASCVD risk factors

- Culturally tailored personalized cardioprotective dietary pattern.
- Emphasize plant-based foods.
- Achieve 5-10% weight loss if overweight.
- Reduce saturated fat: <7% of energy intake.
- Dietary cholesterol: <200 mg/day.
- Reduce added sugars: <10% of energy intake.
- Increase viscous fiber: 5-10 g/day
- Plant sterols/stanols: 2 g/day.

Jacobson et al. J Clin Lipidol. 2015

51



Question

In Outpatient adults, what is the effectiveness and cost benefit of medical nutrition therapy (MNT) provided by an RDN for management of dyslipidemia?

Sikand G et al. J Clin Lipidol October, 2018.

Importance of Culturally Relevant Diet Counseling in SA

- High recidivism rate in subjects who tried to follow eating plans that were different from their usual dietary patterns (Sacks N Eng J Med 2009).
- Culturally tailored cardioprotective dietary pattern and exercise interventions led to improved BMI, A1c, insulin sensitivity and physical activity (Natesan BM/ Open Diabetes Res. Care. 2015).
- Multiple personalized sessions with a dietitian in over 5700 subjects (13 countries) led to improved LDL-C, TG, BMI and A1c (Sikand J Clin Lipidol. 2018).


52


50

Study Selection


- PubMed, MEDLINE, Worldcat.org
- Inclusion criteria:
 - English language
 - Adults over 18
 - MNT provided by a RDN in out patient setting
 - Published Jan 2003-Oct 2014
 - 10 or more subjects in the study with at least 65% completion rate
 - At least one outcome measure of dyslipidemia*


Sikand G et al. J Clin Lipidol October, 2018.






57





56



58

2018 Systematic Review and Meta-analysis Highlights

- Dietitian intervention in 5704 subjects (34 studies) led to improved LDL-C, TG, A1c, BMI, quality adjusted life years and reduced need for lipid-lowering medications.
- Multiple individual sessions with dietitian were clinically and cost beneficial.
- Benefits also reported when dietitian was part of a multidisciplinary health care team.

Sikand et al. J Clin Lipidol. October 2018

Take Away

- Culturally tailored counseling for a personalized cardio-protective dietary pattern.
- Weight loss of 3-5 % of body weight (overweight or obese).
- Reduced intake of saturated fat <7% of energy intake and dietary cholesterol <200 mg/day.
- Reduced intake of added sugars (<10% of total energy).
- 5 to 10 g/day of viscous fiber and 2 g/day of phytosterols.
- Culturally relevant multiple visits with a dietitian for diet and lifestyle counseling.

61

Resources for Practioners

1. Van Horn L et al. Recommended dietary pattern to achieve adherence to the American Heart Association/American College of Cardiology (AHA/ACC) Guidelines: A Scientific Statement from the American Heart Association. *Circulation 2016*;134 e505-e529.

2. Sikand G. "Preventing Heart Disease in Asian Indians: Diet & Lifestyle Recommendations in "Indian Foods: AAPI's (American Association of Physicians from India). Guide to Nutrition, Health and Diabetes" 2011 second edition

3. Palaniappan LP et al. Call to action: CVD in Asian Americans: a science advisory from the American Heart Association. Circulation. 2010;122:1242–1252. doi: 10.1161/CIR.0b013e3181f22af4.

63



- Emphasize a plant based diet with vegetables, fruits, whole grains, legumes, non-tropical vegetable oils, unsalted nuts; non-fat dairy products; and poultry and fish for omnivores.
- Limit sweets, sugar sweetened beverages, fruit juices, salty or highly processed foods and fatty red meat or processed meat.
- Provide culturally appropriate counseling by a health care team. Include a registered dietitian, exercise specialist and stress management counselor.

Jacobson 2015, Vanhorn 2016, Iyengar 2016, Grundy 2018, Arnett 2019

62

NLA Resources for clinicians and patients from NLA


www.lipid.org/CLMT

Follow link to Clinical lifestyle Modification Toolkit (CLMT):

- Heart-Healthy Eating South Asian/ Indian Style
- DASH Dietary Pattern
- Mediterranean style Dietary Pattern
- Vegetarian/Vegan Dietary Pattern
- And many more

certificate.

64





 Remember: If you used your phone to call in, and want CE credit for attending, please send an email with your name to <u>cope@villanova.edu</u> so you receive your

